

2-2021

Assessment of Hyperfibrinolysis in Liver Transplantation Surgery as Measured by Rotational Thromboelastometry

Ryan Madsen

Uzung Yoon

Marc Torjman

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2023_phase1



Part of the [Translational Medical Research Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.



**Sidney Kimmel
Medical College™**
at Thomas Jefferson University

Incidence of Fibrinolysis in Modern Liver Transplantation

Ryan Madsen, Dr. Uzung Yoon*

(*) indicates primary project advisor

Disclosures: None

Acknowledgements: Dr. Uzung Yoon, Paul Del Prato, and Dr. Marc Torjman

Introduction to Liver transplantation fibrinolysis

- Fibrinolysis is a recognized problem during liver transplant surgeries, especially during the reperfusion phase¹
- Modern liver transplantations use significantly less amounts of blood products compared to historical liver transplantations ²
- The rate of fibrinolysis may have changed alongside improving surgical techniques, thus patients may be receiving unnecessary blood products
- Liver transplant surgical techniques may become more efficient, safer, and less costly if fibrinolysis is not happening as frequently as what is currently believed

Research Question

Research question: Using a Rotational thromboelastometry (ROTEM) machine to measure clotting, is the rate of fibrinolysis during modern liver transplant surgery as high as what is currently believed (~15%)?





**Sidney Kimmel
Medical College™**
at Thomas Jefferson University

Hypothesis: The patients who underwent liver transplantation surgeries at Jefferson from 2013-2019 have a significantly lower (<5%) rate of hyperfibrinolysis, as measured by ROTEM clotting parameters, compared to what has been described in previous literature (~15%).


Methods

- Retrospective chart review using ROTEM data from 312 liver transplant surgeries that took place at Jefferson from 2013-2019
 - ROTEM (Rotational thrombelastography): a machine capable of intraoperatively assessing several qualities of a clot from a whole blood assay
 - Rate of fibrinolysis, clot formation time, clot strength etc.
- Data collection: The data was taken from the ROTEM machines and has been matched with patient information from EPIC
- Analysis: Blood was collected at several distinct time points during the surgery and the clotting profile was assessed
- Outcome of interest: Is hyperfibrinolysis* present at any point during the surgery?

*Hyperfibrinolysis is defined a Maximum Lysis Value greater than 15% on NATEM
And a correction of hyperfibrinolysis on APTTEM

Methods continued

- An example of the type of data that ROTEM provides ³
- Note, this information is available during surgery



Results

- Population:
 - Average age at transplantation was 58 ± 11.2 years
 - About 67% of the recipients were men
 - Alcohol (24%), Hepatitis C (22%), and HCC (21%) were the most common reasons for transplantation
- Hyperfibrinolysis:
 - There were 20 patients out of 321 (6.1%) who had hyperfibrinolysis
 - Factors not found to be related to hyperfibrinolysis
 - MELD score
 - INR
 - Intraoperative clot strength
 - Factors found to be related to hyperfibrinolysis
 - Non-alcohol reason for transplantation
 - Multiple causative factors of liver injury
 - Age
 - Third stage of surgery




Conclusions

- Overall, the study appears to show that hyperfibrinolysis occurs less often than what is currently believed
- Potential predicative factors such as age, and reason for transplantation have been identified
- Hyperfibrinolysis remains a serious concern during liver transplantation surgeries and still needs to be rigorously monitored



Future Directions

- Search the data base for additional clotting parameters that can be used to predict coagulopathies
- Gather additional demographic and anesthesia information from the surgeries



References

1. Spiel AO, Mayr FB, Firbas C, Quehenberger P, Jilma B. Validation of rotation thrombelastography in a model of systemic activation of fibrinolysis and coagulation in humans. *J Thromb Haemost*. 2006;4(2):411-416. doi:10.1111/j.1538-7836.2006.01715.x
2. Butler P, Israel L, Nusbacher J, Jenkins DE Jr, Starzl TE. Blood transfusion in liver transplantation. *Transfusion*. 1985;25(2):120–123. doi:10.1046/j.1537-2995.1985.25285169201.x
3. Görlinger, Klaus & Dirkmann, Daniel & Hanke, Alexander. (2016). Rotational thromboelastometry (ROTEM®). 10.1007/978-3-319-28308-1_18.